

LAGRANGIAN SELF-SIMILAR SOLUTIONS IN GRADIENT SHRINKING KÄHLER-RICCI SOLITONS

HIKARU YAMAMOTO

ABSTRACT. In this paper, we give a lower bound estimate for the diameter of a Lagrangian self-shrinker in a gradient shrinking Kähler-Ricci soliton as an analog of a result of A. Futaki, H. Li and X.-D. Li [4] for a self-shrinker in a Euclidean space. We also prove an analog of a result of H.-D. Cao and H. Li [1] about the non-existence of compact self-expanders in a Euclidean space.

1. INTRODUCTION

A gradient shrinking Kähler-Ricci soliton is a Kähler manifold (N, ω, g, J) with a smooth function $f : N \rightarrow \mathbb{R}$ satisfying

$$(1) \quad \text{Ric}(g) + \text{Hess}_g f = g.$$

By the equation (1), it follows that the $(2, 0)$ -part of $\text{Hess} f$ is zero. Hence it is clear that the $(1, 0)$ -part of ∇f is a holomorphic vector field on N . By a simple calculation, it is proved that the gradient of $R(g) + |\nabla f|^2 - 2f$ is zero, and we put a constant C_0 by

$$(2) \quad C_0 := R(g) + |\nabla f|^2 - 2f,$$

where $R(g)$ is the scalar curvature of (N, g) . It is proved that $R(g) \geq 0$ for a complete gradient shrinking Ricci soliton by an application of Corollary 2.5 in [2].

For an immersion $F : L \rightarrow N$, we get a section $(\nabla f) \circ F \in \Gamma(L, F^*(TN))$, and we usually omit the symbol $\circ F$, for short.

Definition 1.1. An immersion map $F : L \rightarrow N$ is called a self-similar solution if it satisfies

$$(3) \quad H = \lambda \nabla f^\perp$$

for some constant $\lambda \in \mathbb{R}$, where H is the mean curvature vector field of F and $\perp$ denotes the projection onto the normal bundle of L . It is called a self-shrinker, a steady soliton or a self-expander when $\lambda < 0$, $\lambda = 0$ or $\lambda > 0$, respectively.

For example, a function $f(z^1, \dots, z^m) := \frac{1}{2}(|z^1|^2 + \dots + |z^m|^2)$ on $\mathbb{C}^m$ with the standard Kähler structure satisfies the identity (1), and it satisfies $\nabla f(x) = x$ under a natural identification of points and tangent vectors for all points $x \in \mathbb{C}^m \cong \mathbb{R}^{2m}$. Hence the equation (3) coincides with $H_x = \lambda x^\perp$ for all points $x \in F(L) \subset \mathbb{C}^m \cong \mathbb{R}^{2m}$, and Definition 1.1 can be considered as a generalization of a self-similar solution in a Euclidean space to in a gradient shrinking Ricci soliton.

2010 *Mathematics Subject Classification.* 53C42, 53C44.

Key words and phrases. mean curvature flow, Ricci flow, self-similar solution, gradient soliton.

This work was supported by Grant-in-Aid for JSPS Fellows Grant Number 13J06407 and the Program for Leading Graduate Schools, MEXT, Japan.

There are many results about self-similar solutions in a Euclidean space. By a generalization of the notion of a self-similar solution in a Euclidean space to in a gradient shrinking Ricci soliton as in Definition 1.1, we can discuss which results about self-similar solutions in a Euclidean space also hold in a gradient shrinking Ricci soliton. As an example of such results, it is proved that a part of a result due to Smoczyk also holds in a gradient shrinking Kähler-Ricci soliton. More precisely, in the proof of Theorem 2.3.5 in [7], Smoczyk proved that every compact Lagrangian self-similar solution with exact mean curvature form is a minimal submanifold in $\mathbb{C}^n$, and as a generalization of this statement, it is proved in [8] that every compact Lagrangian self-similar solution with exact mean curvature form is a minimal submanifold in a gradient shrinking Kähler-Ricci soliton.

In this paper, we give further two results which are already established when (N, g) is a Euclidean space. The first result is an analog of Theorem 4.3 of A. Futaki, H. Li and X.-D. Li [4] under the Lagrangian assumption. This gives a lower bound of the diameter of a Lagrangian self-shrinker in a gradient shrinking Kähler-Ricci soliton.

Theorem 1.2. *Let (N, ω, g, J) be a $2m$ -dimensional gradient shrinking Kähler-Ricci soliton with potential function $f: N \rightarrow \mathbb{R}$ satisfying the equation (1). Let $F: L \rightarrow N$ be a compact Lagrangian self-shrinker with*

$$H = -\frac{1}{2}\nabla f^\perp.$$

Assume that $F(L)$ is not contained in $\{f = m - \frac{C_0}{2}\}$, where C_0 is a constant defined by (2). Then we have

$$\text{diam}(L, F^*g) \geq \frac{\pi}{\sqrt{\frac{3}{4} + \frac{m}{2}(K_0 + A_0^2)}},$$

for constants $K_0, A_0 \geq 0$ satisfying $|K_N| \leq K_0$ and $|A| \leq A_0$, where K_N is the sectional curvature of (N, g) and A is the second fundamental form of F .

The second result is an analog of Proposition 5.3 of H.-D. Cao and H. Li [1] under the Lagrangian assumption. This theorem states the non-existence of compact Lagrangian self-expanders in a certain gradient shrinking Kähler-Ricci soliton.

Theorem 1.3. *Let (N, ω, g, J) be a $2m$ -dimensional gradient shrinking Kähler-Ricci soliton with potential function $f: N \rightarrow \mathbb{R}$ satisfying the equation (1) and assume that $R(g) < 2m$, then there exists no compact Lagrangian self-expander in N .*

The rest of this paper is organized as follows. In Section 2, we give some characterization of self-similar solutions in gradient shrinking Ricci solitons. In Section 3, we give a proof of Theorem 1.2 and 1.3.

Acknowledgements. I would like to thank my supervisor, A. Futaki for many useful discussions and constant encouragement.

2. CHARACTERIZATION OF SELF-SIMILAR SOLUTIONS

In this section, we give some characterization of self-similar solutions in gradient shrinking Ricci solitons and review a result in [8].

The first characterization is given by the variation of the weighted volume as follows. Let (N, g, f) be a n -dimensional gradient shrinking Ricci soliton with

potential function f satisfying (1). For an m -dimensional compact manifold L and a constant $\lambda \in \mathbb{R}$, we define the weighted volume functional $\mathcal{F}_\lambda$ by

$$\mathcal{F}_\lambda(F) := \int_L e^{\lambda f} d\mu(F^*g)$$

for each immersion $F: L \rightarrow N$, where $d\mu(F^*g)$ is the induced measure on L with respect to the metric F^*g .

Proposition 2.1. *Let $F: L \rightarrow N$ be an immersion and $\lambda \in \mathbb{R}$ be a constant. Then the following three conditions are equivalent.*

- (1) F is a self-similar solution with $H = \lambda \nabla f^\perp$.
- (2) F is a minimal immersion with respect to a metric $e^{2\lambda f/m}g$ on N .
- (3) F is a critical point of $\mathcal{F}_\lambda$.

The equivalence of (1) and (2) is proved in [8], and the equivalence of (2) and (3) can be easily proved by the equality

$$\int_L e^{\lambda f} d\mu(F^*g) = \int_L d\mu(F^*(e^{2\lambda f/m}g)).$$

The equivalence of (1) and (3) can be considered as a generalization of Proposition 3.6 in [3].

The second characterization is given by the asymptotic behavior of a Ricci-mean curvature flow, the coupled equation of the Ricci flow and the mean curvature flow. Let (N, ω, g, J) be a compact $2m$ -dimensional complete gradient shrinking Kähler-Ricci soliton with a potential function $f: N \rightarrow \mathbb{R}$ satisfying (1). Fix a time $T > 0$. Then for $t \in [0, T)$ we define $g_t := (T - t)\Phi_t^*g$, where $\{\Phi_t: N \rightarrow N\}_{t \in (-\infty, T)}$ is the 1-parameter family of holomorphic automorphisms of (N, J) with $\Phi_0 = \text{id}_N$ generated by the time dependent vector field $\frac{1}{2(T-t)}\nabla f$. Then g_t is a solution of Kähler-Ricci flow, that is, the associated Kähler form $\omega_t(\cdot, \cdot) := g_t(J\cdot, \cdot)$ satisfies

$$\frac{\partial}{\partial t}\omega_t = -\rho(\omega_t),$$

where $\rho(\cdot, \cdot) := \text{Ric}(J\cdot, \cdot)$ is the Ricci form of ω_t . Here we review the main result in [8]. Let L be an m -dimensional compact manifold and $F: L \times [0, T) \rightarrow N$ be a solution of Ricci-mean curvature flow along $g_t = (T - t)\Phi_t^*g$, that is, F satisfies

$$\frac{\partial}{\partial t}F_t = H(F_t),$$

where $H(F_t)$ is the mean curvature vector field of $F_t(\cdot) := F(\cdot, t)$ calculated by g_t at each time t . Assume that the initial immersion $F_0: L \rightarrow N$ is a Lagrangian immersion for the initial Kähler form ω . Then, it follows that $F_t: L \rightarrow N$ is also a Lagrangian immersion with respect to ω_t for all $t \in [0, T)$ (c.f. [6]). That is, the Lagrangian condition is preserved under a Ricci-mean curvature flow along a Kähler-Ricci flow. We further assume that F develops a singularity of type I, that is, the norm of the second fundamental form of F_t (denoted by $A(F_t)$) satisfies

$$\limsup_{t \rightarrow T} (\sqrt{T - t} \max_L |A(F_t)|) < \infty.$$

Then, in [8], it is proved that for any sequence $t_j \rightarrow T$ and any point $p_0 \in L$ the family of pointed immersions $\tilde{F}_j: (L, p_0) \rightarrow N$ defined by $\tilde{F}_j := \Phi_{t_j} \circ F_{t_j}$

subconverges to a pointed immersion $\tilde{F}_\infty: (L_\infty, p_\infty) \rightarrow N$ satisfying

$$H(\tilde{F}_\infty) = -\frac{1}{2}\nabla f^\perp.$$

This result can be considered as a generalization of Huisken's result in [5] for a mean curvature flow in a Euclidean space.

Since each $\tilde{F}_j$ is a Lagrangian immersion in (N, ω) and the Lagrangian condition $(\tilde{F}_j^* \omega = 0)$ is a closed condition, it follows that $\tilde{F}_\infty: L_\infty \rightarrow N$ is a Lagrangian immersion, that is, the Lagrangian self-shrinker. Hence a Lagrangian self-shrinker is an asymptotic model of a Lagrangian mean curvature flow with a type I singularity along a Kähler-Ricci flow constructed from a gradient shrinking Kähler-Ricci soliton.

Remark 2.2. Actually, the same statement also holds under some additional assumptions even though N is non-compact and complete, see [8] for detail. The differences of factor 2 or 1/2 between coefficients appeared in some formula in this paper and those in [8] arise from the difference of factor 2 between the Ricci flow equation $\partial_t g_t = -2\text{Ric}(g_t)$ and the Kähler-Ricci flow equation $\partial_t \omega_t = -\rho(\omega_t)$.

3. PROOFS OF THEOREM 1.2 AND 1.3

First, we give a proof of Theorem 1.2. The proof is an analog of the proof of Theorem 4.3 of [4]. As the first step, we prove that the weighted Laplacian $\underline{\Delta}_\phi$ defined below has an eigenvalue 1. In the second step, we use Theorem 1.1 in [4] giving an estimate for the first eigenvalue of the weighted Laplacian.

Proof of Theorem 1.2. Put a smooth function on L by $\phi := \frac{1}{2}f \circ F$ and consider the weighted Laplacian

$$\mathcal{L} := \underline{\Delta}_\phi := \underline{\Delta} - \bar{\nabla} \phi \cdot \bar{\nabla}$$

which acts on $C^\infty(L)$, where $\underline{\Delta}$ and $\bar{\nabla}$ is the Laplacian and the gradient on (L, F^*g) . Since

$$(4) \quad \underline{\Delta}(f \circ F) = \text{tr}^\top \text{Hess}_g f + g(\nabla f, H),$$

we have

$$\begin{aligned} \mathcal{L}\phi &= \frac{1}{2}\underline{\Delta}(f \circ F) - \frac{1}{4}(F^*g)(\bar{\nabla}(f \circ F), \bar{\nabla}(f \circ F)) \\ &= \frac{1}{2}(\text{tr}^\top \text{Hess}_g f + g(\nabla f, H)) - \frac{1}{4}|\nabla f^\top|^2, \end{aligned}$$

where $\text{tr}^\top$ is the tangential trace, that is, $\text{tr}^\top B := \text{tr}_{F^*g}(F^*B)$ for a 2-tensor B on N . Since $H = -\frac{1}{2}\nabla f^\perp$, we have

$$\frac{1}{2}g(\nabla f, H) - \frac{1}{4}|\nabla f^\top|^2 = -\frac{1}{4}|\nabla f^\perp|^2 - \frac{1}{4}|\nabla f^\top|^2 = -\frac{1}{4}|\nabla f|^2.$$

Since (N, g) is a gradient shrinking Ricci soliton satisfying (1), we have

$$(5) \quad \text{tr}^\top \text{Hess}_g f = \text{tr}^\top (g - \text{Ric}(g)) = m - \text{tr}^\top \text{Ric}(g).$$

Furthermore, since $F : L \rightarrow N$ is a Lagrangian immersion and (N, g, J) is a Kähler manifold, we have

$$(6) \quad \begin{aligned} \operatorname{tr}^\top \operatorname{Ric}(g) &= \sum_{i=1}^m \operatorname{Ric}(g)(F_* e_i, F_* e_i) \\ &= \frac{1}{2} \sum_{i=1}^m \operatorname{Ric}(g)(F_* e_i, F_* e_i) + \frac{1}{2} \sum_{i=1}^m \operatorname{Ric}(g)(JF_* e_i, JF_* e_i) = \frac{1}{2} R(g), \end{aligned}$$

for an orthonormal basis $e_1, \dots, e_m$ on (L, F^*g) . Hence we have

$$\mathcal{L}\phi = \frac{m}{2} - \frac{1}{4}(R(g) + |\nabla f|^2) = \frac{m}{2} - \frac{1}{4}(C_0 + 2f) = \left(\frac{m}{2} - \frac{C_0}{4}\right) - \phi.$$

Since $\mathcal{L}(\text{const}) = 0$, we have

$$\mathcal{L}\left(\left(\frac{m}{2} - \frac{C_0}{4}\right) - \phi\right) = -\left(\left(\frac{m}{2} - \frac{C_0}{4}\right) - \phi\right).$$

By the assumption, $\frac{m}{2} - \frac{C_0}{4} - \phi \neq 0$. Hence we have proved that 1 is an eigenvalue of the weighted Laplacian $\mathcal{L} = \Delta_\phi$.

Next, we prove that

$$(7) \quad \operatorname{Ric}(F^*g) + \operatorname{Hess}_{F^*g}\phi \geq \kappa F^*g,$$

for a Riemannian manifold (L, F^*g) , where

$$\kappa := \frac{1}{2} - m(K_0 + A_0^2).$$

Let X be a tangent vector on L and $e_1, \dots, e_m$ be an orthonormal basis on (L, F^*g) . Then, by the Gauss equation, we have

$$\begin{aligned} \operatorname{Ric}(F^*g)(X, X) &= \sum_{i=1}^m \operatorname{Rm}(F^*g)(X, e_i, X, e_i) \\ &= \sum_{i=1}^m \operatorname{Rm}(g)(F_*X, F_*e_i, F_*X, F_*e_i) \\ &\quad - \sum_{i=1}^m |A(X, e_i)|^2 + g(A(X, X), H). \end{aligned}$$

Furthermore, we have

$$\begin{aligned} \operatorname{Hess}_{F^*g}\phi(X, X) &= \frac{1}{2} \operatorname{Hess}_{F^*g}(f \circ F)(X, X) \\ &= \frac{1}{2} \left(\operatorname{Hess}_g f(F_*X, F_*X) + g(A(X, X), \nabla f) \right) \\ &= \frac{1}{2} |X|^2 - \frac{1}{2} \operatorname{Ric}(g)(F_*X, F_*X) - g(A(X, X), H), \end{aligned}$$

where we used $\text{Hess}_g f = g - \text{Ric}$ and $\nabla f^\perp = -2H$. Since

$$\begin{aligned} & \sum_{i=1}^m \text{Rm}(g)(F_*X, F_*e_i, F_*X, F_*e_i) - \frac{1}{2} \text{Ric}(g)(F_*X, F_*X) \\ &= \frac{1}{2} \sum_{i=1}^m \text{Rm}(g)(F_*X, F_*e_i, F_*X, F_*e_i) - \frac{1}{2} \sum_{i=1}^m \text{Rm}(g)(F_*X, JF_*e_i, F_*X, JF_*e_i) \\ &\geq -\frac{1}{2}mK_0|X|^2 - \frac{1}{2}mK_0|X|^2 = -mK_0|X|^2 \end{aligned}$$

and

$$-\sum_{i=1}^m |A(X, e_i)|^2 \geq -mA_0|X|^2,$$

we have

$$\text{Ric}(F^*g)(X, X) + \text{Hess}_{F^*g}\phi(X, X) \geq \left(\frac{1}{2} - m(K_0 + A_0^2)\right) |X|^2.$$

Hence the inequality (7) holds.

Thus, by Theorem 1.1 in [4], we have

$$(8) \quad 1 \geq \sup_{s \in (0,1)} \left\{ 4s(1-s) \frac{\pi^2}{d^2} + s\kappa \right\},$$

where $d := \text{diam}(L, F_*g)$. Choosing $s = \frac{1}{2}$ in (8), we have

$$d \geq \frac{\pi}{\sqrt{1 - \frac{1}{2}\kappa}} = \frac{\pi}{\sqrt{\frac{3}{4} + \frac{m}{2}(K_0 + A_0^2)}}.$$

Hence the proof is completed. $\square$

Next, we prove Theorem 1.3. The proof is an analog of the proof of Proposition 5.3 of [1].

Proof of Theorem 1.3. Assume that there exists a compact Lagrangian self-expander $F: L^m \rightarrow N^{2m}$ (*compact* means M is compact) with $H = \lambda \nabla f^\perp$ for some positive constant $\lambda > 0$. Then, using computations (4), (5) and (6) in the proof of Theorem 1.2 and the equality $H = \lambda \nabla f^\perp$, we have

$$\begin{aligned} \underline{\Delta}(f \circ F) &= \text{tr}^\top \text{Hess}_g f + g(\nabla f, H) \\ &= m - \frac{1}{2}R(g) + \frac{1}{\lambda}|H|^2, \end{aligned}$$

where $\underline{\Delta}$ is the Laplacian on (L, F^*g) . Hence, by the divergence theorem, we have

$$0 = \int_L \underline{\Delta}(f \circ F) d\mu(F^*g) = \int_L \left(m - \frac{1}{2}R(g) + \frac{1}{\lambda}|H|^2 \right) d\mu(F^*g).$$

However, the right hand side is strictly positive by the assumption $R(g) < 2m$. This leads to a contradiction. $\square$

REFERENCES

- [1] H.-D. Cao and H. Li. A gap theorem for self-shrinkers of the mean curvature flow in arbitrary codimension. *Calc. Var. Partial Differential Equations* 46 (2013), no. 3-4, 879–889.
- [2] B.-L. Chen. Strong uniqueness of the Ricci flow. *J. Differential Geom.*, 82 (2009), no. 2, 363–382.
- [3] T. H. Colding and W. P. Minicozzi II. Generic mean curvature flow I: generic singularities. *Ann. of Math. (2)* 175 (2012), no. 2, 755–833.
- [4] A. Futaki, H. Li and X.-D. Li. On the first eigenvalue of the Witten-Laplacian and the diameter of compact shrinking solitons. *Ann. Global Anal. Geom.* 44 (2013), no. 2, 105–114.
- [5] G. Huisken. Asymptotic behavior for singularities of the mean curvature flow. *J. Differential Geom.*, 31(1):285–299, 1990.
- [6] J. D. Lotay and T. Pacini. Coupled flows, convexity and calibrations: Lagrangian and totally real geometry. arXiv:1404.4227.
- [7] K. Smoczyk. The Lagrangian mean curvature flow. Univ. Leipzig (Habil.-Schr.), 2000.
- [8] H. Yamamoto. Ricci-mean curvature flow in gradient shrinking Ricci solitons. arXiv:1501.06256.

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE, TOKYO UNIVERSITY OF SCIENCE, 1-3
KAGURAZAKA, SHINJUKU-KU, TOKYO 162-8601, JAPAN
E-mail address: hyamamoto@rs.tus.ac.jp